

Nutritional, Physiological and Perceptual Responses During a Summer Ultra-Endurance  
Cycling Event

Lawrence E. Armstrong,  
Douglas J. Casa,  
Holly Emmanuel,  
Matthew S. Ganio,  
Jennifer F. Klau,  
Elaine C. Lee,  
Carl M. Maresh,  
Brendon P. McDermott,  
Rebecca L. Stearns,  
Jakob L. Vingren,  
Jonathan E. Wingo,  
Keith H. Williamson,  
and  
Linda M. Yamamoto

Deposited 12/18/2017

Citation of published version:

Armstrong, L. E., Casa, D. J., Emmanuel, H., Ganio, M. S., Klau, J. F., Lee, E. C., & ... Yamamoto, L. M. (2012). Nutritional, physiological, and perceptual responses during a summer ultraendurance cycling event. *Journal Of Strength And Conditioning Research*, (2), 307. doi: 10.1519/JSC.0b013e318240f677

Available at: [http://journals.lww.com/nsca-jscr/Abstract/2012/02000/Nutritional,\\_Physiological,\\_and\\_Perceptual.1.aspx](http://journals.lww.com/nsca-jscr/Abstract/2012/02000/Nutritional,_Physiological,_and_Perceptual.1.aspx)

**Nutritional, Physiological and Perceptual Responses  
During a Summer Ultra-Endurance Cycling Event**

First Author & Corresponding Author: Lawrence E. Armstrong

Manuscript R-336911

**Authors:** Lawrence E. Armstrong, Douglas J. Casa, Holly Emmanuel,  
Matthew S. Ganio, Jennifer F. Klau, Elaine C. Lee, Carl M. Maresh  
Brendon P. McDermott, Rebecca L. Stearns, Jakob L. Vingren,  
Jonathan E. Wingo, Keith H. Williamson, Linda M. Yamamoto

**Institution:** University of Connecticut, Human Performance Laboratory,  
Dept. of Kinesiology, Storrs, CT 06269-1110, USA

**Corresponding Author:** Lawrence E. Armstrong  
Tel: 860 486 2647; FAX: 860 486 1123  
e-mail: [lawrence.armstrong@uconn.edu](mailto:lawrence.armstrong@uconn.edu)

**Running Head:** Responses to Summer Ultra-Endurance Cycling

[7 Tables, 4 Figures]

This research was not funded via external sources. All funds were internal to the university.

**Nutritional, Physiological and Perceptual Responses  
During a Summer Ultra-Endurance Cycling Event**

**Running Head:** Responses During Ultra-Endurance Cycling

[7 Tables, 4 Figures]

## ABSTRACT

Despite the rapid growth of mass-participation road cycling, little is known about the dietary, metabolic and behavioral responses of ultra-endurance cyclists. This investigation describes physiological responses, perceptual ratings, energy balance, and macronutrient intake of 42 men (mean  $\pm$  SD; age,  $38\pm 6$  y; height,  $179.7\pm 7.1$  cm; body mass,  $85.85\pm 14.79$  kg) and 6 women (age,  $41\pm 4$  y; height,  $168.0\pm 2.9$  cm; body mass,  $67.32\pm 7.21$  kg) during a summer 164-km road cycling event. Measurements were recorded one day before, and on Event Day (10.5 h) at the start (0 km), at two aid stations (52 km and 97 km), and at the finish line (164 km). Ambient temperature was  $>39.0^{\circ}\text{C}$  during the final 2 h of exercise. Mean finish times for men ( $9.1\pm 1.2$  h) and women ( $9.0\pm 0.2$  h) were similar, as were mean gastrointestinal temperature [ $T_{\text{GI}}$ ], four hydration biomarkers, and five perceptual (e.g., thermal, thirst, pain) ratings. Male cyclists consumed enough fluids on Event Day ( $5.91\pm 2.38$  L; 49% water) to maintain body mass within 0.76 kg, start to finish, despite a sweat loss of  $1.13\pm 0.54$  L $\cdot$ h $^{-1}$  and calculated energy expenditure of  $3,115$  Kcal $\cdot$ 10.5h $^{-1}$ . However, men voluntarily underconsumed food energy (deficit of 2,594 Kcal, 10.9 MJ), and specific macronutrients (carbohydrates,  $106\pm 48$  g; protein,  $8 \pm 7$  g; and sodium,  $852 \pm 531$  mg) between 0530 and 1400 h. Also, a few men exhibited extreme final values (i.e., urine specific gravity of 1.035 to 1.038,  $n=5$ ; body mass loss  $>4$  kg,  $n=2$ ;  $T_{\text{GI}}$ , 39.4 and 40.2). We concluded that these findings provide information regarding energy consumption, macronutrient intake, hydration status and the physiological stresses that are unique to ultra-endurance exercise in a hot environment.

(268 words)

Key words: fluids, electrolytes, urine, sport nutrition, thermoregulation, energy expenditure

## **INTRODUCTION**

The foods and fluids consumed by endurance road cyclists are related to both performance and well-being. Before, during and after competition, dietary strategies can reduce fatigue and the risk of illness, as well as enhance recovery and optimize physiological adaptations (9). Thus, several previous publications have evaluated the dietary intakes of collegiate and Olympic cyclists relative to the metabolic requirements imposed by elite competition (9,15,26). These publications focused on carbohydrate intake before and during exercise because hypoglycemia influences central fatigue (6), endurance is enhanced by pre-exercise glycogen levels, and because exhaustion from glycogen depletion can be postponed by carbohydrate feeding during exercise (19). A large body of information also indicates that dehydration can degrade endurance exercise performance when more than 2% of body mass is lost (12,28). That is, when dehydration exceeds 3-5% of body mass, both sweat rate and skin blood flow decrease, and the risk of exertional heat illness increases (2). Further, human hyperthermia and high ambient temperature potentiate the interactions between exercise metabolism, exercise performance, hydration state, and health. Although few core body temperature measurements in cyclists have been published, we know that marathon runners experience rectal temperatures of 39.0 – 41.7°C after running in mild (7 – 18°C)

environments (21,22). Unfortunately, the scientific literature is limited as to how well non-elite cyclists maintain carbohydrate intake, fluid balance, and body temperature during ultra-endurance events.

During the past decade, the popularity of mass participation endurance cycling events has grown rapidly in many countries. USA Cycling, the official governing body for all disciplines of competitive cycling in the United States, reported 66,844 licensed members in 2009 and sanctioned 2,638 events (31). That year, the Hotter 'n Hell Hundred (HHH) was the largest single-day 100-mile (164 km) event in the United States. Our research team recognized that this endurance event presented unique nutritional, physiological and behavioral stresses to entrants, as well as difficult clinical care challenges to medical personnel, which have not been described for non-elite cyclists.

Therefore, the purpose of the present investigation was to describe the physiological and metabolic responses, perceptual ratings, energy balance, and macronutrient intake of 48 recreational endurance cyclists during a 164-km event conducted in a mean ambient temperature of  $34.4 \pm 4.9^\circ\text{C}$ . We hypothesized that these men and women would not maintain thermal, fluid and energy balance during this 9-h effort, and that behavioral and physiological measures would indicate great stress. We anticipated that the findings of this research would provide information to dietitians, clinicians, coaches and race directors regarding energy consumption, macronutrient intake, hydration status and the physiological stresses that are unique to ultra-endurance exercise in a hot environment.

## METHODS

With the approval of the event directors, investigators recruited cyclists as they attended the registration and pre-race exposition of the HHH event (Day -1). This event occurred in Texas during the month of August. Prior to giving informed written consent, each cyclist received a written and verbal description of all procedures, measurements, time commitment, benefits and risks, as approved by the university Institutional Review Board for Human Studies. Subjects were not paid, but received an explanation of their own data.

Subjects completed a medical history questionnaire and a 28-day exercise history, which subsequently was screened by the race medical director and the responsible investigator. Exclusionary criteria included inadequacy of recent training, present musculo-skeletal injury, and a history of either exertional heatstroke, exercise-heat intolerance, disease of the gastrointestinal tract (e.g., diverticulitis, hypomotility, inflammatory bowel disease), or gastrointestinal surgery. All test participants had previously completed at least one 160-km cycling event and were aged 19-39 years.

Investigators did not provide food or water to participants. Investigators also did not offer advice/instructions to participants about planning or execution of race strategies, or about food and fluid intake.

**Physiological Variables.** On Day -1, the height of each test subject was measured by standing against a tape measure, attached to a wall. Body mass was measured with a floor scale, accurate to  $\pm 100$  g. Body mass index (BMI) was calculated as body mass (kg) divided by height<sup>2</sup> (m<sup>2</sup>). Age was recorded to the nearest year. The characteristics (mean  $\pm$  SD, range) of 42 men were: age, 38 $\pm$ 6 (20-50) y;

height,  $179.7 \pm 7.1$  (163.4-194.0) cm; body mass,  $85.85 \pm 14.79$  (57.6-115.7) kg; BMI,  $26.4 \pm 4.0$  (19.4-37.4). The characteristics of 6 women were: age,  $41 \pm 4$  (36-48) y; height,  $168.0 \pm 2.9$  (164.5-171.3) cm; body mass,  $67.32 \pm 7.21$  kg (58.0-77.6); BMI,  $23.8 \pm 2.3$  (20.9-26.9).

Heart rate (n=16 men, n=3 women) and gastrointestinal temperature ( $T_{GI}$ ; n=12 men, n=3 women) were measured in a subset of participants who were identified at the test subject information meeting. Heart rate was assessed via wrist cardiometer that read heart rhythm from a chest strap. Gastrointestinal temperature ( $T_{GI}$ ) was measured via ingestible thermistor (CorTemp®, HQ Inc., Palmetto, FL), which each volunteer swallowed at 2100 h on the night prior to the event; ingestion was verified by a telephone call from an investigator. By the next morning, this temperature sensor had moved into the digestive tract via peristalsis.  $T_{GI}$  was measured with a hand-held digital thermometer ( $\pm 0.1^\circ\text{C}$ ), positioned near the posterior lumbar curve.

Two days prior to Event Day, a subset of 10 men were identified for a preliminary procedure that allowed calculation of sweat rate (L/min). This procedure has been described in detail elsewhere (1); it involved  $41 \pm 7$  min of outdoor exercise in a hot environment, pre- and post-exercise body mass measurements, and corrections for the volume of sweat trapped in clothing, and the mass of urine excreted. No fluid or food was consumed during this procedure.

Sweat rate also was assessed in 20 males on Event Day; these individuals were selected from a larger sample of riders because they voluntarily consumed no solid food, up to the 52 km point. This calculation involved the body mass difference (0 km to

52 km) with corrections for the mass of fluids consumed, as reported in diet records written at the 52 km aid station.

Energy expenditure during exercise was calculated using an over-ground cycling method (29) that incorporated the ground speed and body mass of each cyclist to derive the rate of oxygen consumption ( $L \cdot \text{min}^{-1}$ ). Considering the elapsed time, oxygen consumption was then converted to Kcal. Metabolic water production, an additional component of total body water balance, was calculated with the method of Hoyt & Honig (17). This technique utilized total energy expenditure ( $\text{kcal} \cdot 24\text{h}^{-1}$ ) including exercise, sleep and other daily activities.

**Perceptual Ratings.** Four rating scales, all relevant to prolonged endurance exercise in a hot environment, were recorded for men and women at the starting line, at two aid stations, and at the finish line. Firstly, the rating of thermal sensation (32) included seventeen categories of thermal comfort, ranging from 0.0 (“unbearably cold”) to 8.0 (“unbearably hot”), in increments of 0.5 unit. Secondly, a rating of thirst (13) included nine categories ranging from 1 (“not thirsty at all”) to 9 (“very, very thirsty”). Thirdly, a 6 to 20 point rating of perceived exertion (5) was presented to each cyclist after exercise, at the two aid stations and at the finish line; the extreme category terms ranged from “very, very light” to “very, very hard”. Cyclists also responded to the statement “I have a muscle cramp” by selecting one of the following options: not at all, a little, somewhat, moderate, a lot, extreme; these options were scored 1 through 6, for statistical analysis. Finally, the pain rating scale (11) involved twelve categories, with extreme ratings of 0 (“no pain at all”) to 10 (“Extremely intense pain, almost unbearable”).

**Diet Records.** During the 24 h prior to this event (Day -1), cyclists recorded all food and fluid that he/she consumed during meals and snacks. These records followed written and verbal instructions, provided by an investigator at the pre-participation briefing meeting. Cyclists recorded details such as the number, volume, size, brand, manufacturer, and method of preparation; nutrition labels and packages were submitted, when possible. The Day -1 diet records were reviewed for completeness by an investigator, in the presence of each cyclist, on the morning of the HHH.

Diet records also were maintained on the Event Day; these included the morning meal, snacks and all food-fluid items consumed during the event. At two aid stations (52 km and 97 km) and at the finish line (164 km), an investigator examined diet records for completeness, verified individual items, and reminded cyclists to record all items which had been consumed.

Dietary records from Day -1 and Event Day were analyzed by selecting individual food items from a commercial data base (Nutritionist Pro™, version 1.2, N-Squared Computing, Salem, OR). Computed outputs included the macronutrients and fluids that appear below.

**Data Measurement Sites.** On Day -1, each participant received written and verbal instructions regarding the procedures and measurements that would occur at four data collection points. Measurements were taken at the same four locations for all cyclists: at the main medical tent (0 km, prior to the event), at two aid stations on the course (52 km and at 97 km), and at the finish line (164 km) within the same medical tent utilized prior to the race start. Data sheets and rating scales were identical at all sites.

On Event Day prior to the 0700 h start, subjects reported to a medical tent near the starting line, located in the center of Wichita Falls, Texas. Baseline measurements of body mass and perceptual ratings were recorded. Floor scales (Ohaus, model DS44L, Florham Park, NJ) were calibrated to a tolerance of  $\pm 100\text{g}$ . An investigator verified reception of a signal from the transmitter-thermistor that had been swallowed on the previous night, and the initial pre-exercise  $T_{GI}$  was measured. A urine sample was collected and analyzed for urine specific gravity (hand-held refractometer; model 300CL; Atago Co., Tokyo, Japan) and urine color (3). A single blood sample was collected from an antecubital vein, centrifuged, and analyzed in triplicate for osmolality with a freezing point osmometer (Advanced Instruments, Needham Heights, MA, model 3250). An aliquot of blood was frozen for later analysis in triplicate of the serum electrolytes sodium, chloride and potassium (ion selective electrodes; Medica Corp., Bedford, MA). Both the osmometer and the electrolyte analyzer were calibrated using standard solutions and methods provided by the manufacturer. An investigator then gave participants a verbal description of the physiological and perceptual measurements that would be taken that day at each data collection site.

At the aid stations (52 and 97 km, located on rural roads surrounding Wichita Falls, TX), heart rate and  $T_{GI}$  were recorded for selected cyclists, immediately after exercise. For all participants, body mass (wearing the same clothing worn at the starting line), urine specific gravity, urine color, and five perceptual ratings were measured. Body weight scales at all locations were newly purchased and manufactured by the same company. All scales were calibrated prior to the event,

using weights of known mass. Identical refractometers were utilized at each site; each was calibrated prior to the event, using distilled water, per manufacturer instructions.

Cyclists were asked to remember all food and fluid which they consumed during the next event stage, immediately before they mounted bicycles and proceeded toward the next data collection site.

After completing the entire 164 km distance, cyclists reported to the medical tent near the finish line. The heart rates and  $T_{GI}$  of selected subjects were measured immediately after exercise. Blood and urine were sampled and analyzed as described above. All other physiological and perceptual variables which had been measured previously at the starting line (see above), including diet records, were repeated.

**Statistical Analyses.** All data are presented as means  $\pm$  SD. Repeated measure analysis of variance (ANOVA) were applied to all variables that were measured at the starting line, the 52 km aid station, the 97 km aid station, and finish line. A t-test for paired samples was used to compare Pre- and Post-event means, as well as Day -1 and Event Day means, Although the authors believe that separating the data on the basis of sex is informative and unique, no statistical comparisons of male and female data (across time or between sexes) were performed due to the small number of women who participated.

The number of test subjects varies in the following tables and figures, for the following reasons: investigators could not locate all subjects at field measurement sites because thousands of cyclists participated, test subjects chose not to report certain findings, and test subject attrition.

## RESULTS

Environmental conditions on the Event Day were recorded each hour, from 0800 to 1500 h. The mean ( $\pm$  SD) dry bulb temperature was  $34.4 \pm 5.0^\circ\text{C}$ , ranging from  $24.4$  (0800 h) to  $39.5^\circ\text{C}$  (1400 h). The relative humidity was  $53 \pm 14$ ; it ranged from  $40$  (1400 h) to  $83\%$  (0800 h). The wet bulb globe temperature was  $30.8 \pm 3.5^\circ\text{C}$ , and ranged from  $22.8$  (0800 h) to  $33.6^\circ\text{C}$  (1400 h). Cloud cover throughout Event Day was minimal or nonexistent.

Registered entrants included 704 men and women, and 96.2% finished the entire 164 km distance. The mean finish time for men and women was similar (Table 1); also, the times to reach the 52 km and 97 km aid stations were similar. We estimated that these times included 10 min for rest, rehydration, and data collection at each aid station. The average finishing times of men (9.1 h) and women (9.0 h) demonstrate that subjects were not elite competitors, in that the times of the fastest finishers were as follows: men, 93 cyclists finished in  $<5$  h, 194 cyclists finished between 5-6 h, and 163 cyclists finished between 6-7 h; women, 5 cyclists finished in  $<5$  h, 10 cyclists finished between 5-6 h, and 24 cyclists finished between 6-7 h. The range of speeds and finish times were  $14.3$ - $22.7$   $\text{km}\cdot\text{h}^{-1}$  and 7.1-11.3 h for men, and  $17.5$ - $18.5$   $\text{km}\cdot\text{h}^{-1}$  and 8.7-9.2 h for women.

The mean final heart rate for men significantly increased ( $P < .001$ ) between the 52 km mark and the finish line (Table 1). The mean ( $\pm$  SD) gastrointestinal temperature (Table 1) increased significantly ( $P < .05$  to  $.001$ ) in 12 men to  $38.5^\circ\text{C}$ ; it increased to  $38.1^\circ\text{C}$  in women ( $n=3$ ). These mean values suggest only mild hyperthermia.

However, the frequency distribution of  $T_{GI}$  (Figure 1) shows that considerable variability existed, and that three cyclists experienced a  $T_{GI}$  increase of  $>2.0^{\circ}\text{C}$ .

Unexpectedly, five men reported to investigators, before starting their 160-km ride, with early morning urine specific gravities of 1.030 to 1.037, indicating frank dehydration. As anticipated, both mean urine specific gravity and mean urine color indicated progressive dehydration during the event (Table 2), but the final (164 km) mean values represented a wide range of hydration categories (4). For example, the frequency distribution of final urine specific gravities (Figure 2) showed that 11 men experienced a final urine specific gravity  $>1.030$ , and that two finished the event with  $<1.010$ . Urine color was strongly and positively correlated with urine specific gravity ( $r^2 = 0.86$ ;  $P < .0001$ ), verifying previously published findings (3), and suggesting a simple option for personal hydration assessment.

As noted above, sweat rate was measured on two different occasions. Two days prior to Event Day (Thursday), a mean sweat rate of  $1.41 \pm 0.51 \text{ L}\cdot\text{h}^{-1}$  (41 min of exercise) was measured for a group of 10 men. On the Event Day (Saturday), the mean sweat rate was  $1.13 \pm 0.54 \text{ L}\cdot\text{h}^{-1}$  for twenty men during the first 52 km (elapsed time, 1.8 h; mean  $T_{GI}$ ,  $38.14^{\circ}\text{C}$ ) of the event. Our casual observations suggested that exercise intensity on Thursday was uniformly greater than on Event Day and that air temperatures were similar; this likely explained the difference in sweat rate. Although few field measurements of sweat rate have been reported for cyclists who exercise in hot environments, these values are comparable to male sweat rates recorded during summer training sessions for soccer and rowing (28).

Mean data in Table 2 suggest that both men and women maintained body mass well during the 164 km ride. Men lost an average of 0.76 kg (0.9% of body mass) and women lost only 0.07 kg (0.1% of body mass). Because it was logistically impossible to measure the mass of food items (i.e., with cyclists moving independently along a 164 km course), we could not determine the extent to which body mass was offset by solid foods; this is acknowledged as a limitation of body mass data. However, the frequency distribution of body mass change (Figure 3) shows that considerable variability existed, and that two men finished the event with a loss of more than 4.0 kg; this represents a considerable dehydration. Seventeen cyclists gained body mass (0.1 – 3.0%; Fig. 3), but we do not know if this was due to solid food or fluid retention.

Mean serum osmolality and serum concentrations of sodium and chloride (Table 3) were statistically similar, before and after exercise. Serum potassium increased significantly in men ( $P < .001$ ) during 9 h of exercise; this likely was due to the release of intramuscular potassium into the circulation. The range of serum values at the finish line (164 km) for men ( $n = 33$ ) were as follows: osmolality, 280–304 mOsm·kg<sup>-1</sup>; sodium, 134–148 mEq·L<sup>-1</sup>; chloride, 94–109 mEq·L<sup>-1</sup>; and potassium, 3.6–5.5 mEq·L<sup>-1</sup>. These ranges are similar to published clinical norms of a large sample of healthy individuals (24): osmolality, 285–295 mOsm·kg<sup>-1</sup>; sodium, 136–145 mEq·L<sup>-1</sup>; chloride, 98–106 mEq·L<sup>-1</sup>; and potassium, 3.5–5.0 mEq·L<sup>-1</sup>.

Five perceptual ratings, all relevant to endurance exercise, are reported in Table 4. Similar to the physiological responses above, the mean perceptual ratings indicated mild-to-moderate strain. As anticipated, thermal sensations, thirst, perceived exertion, pain and sensation of muscle cramping increased significantly ( $P < .025$  to  $.001$ ) from the

start (0 km) to the finish (164 km), but the mean values demonstrated relatively small changes. The range of perceptual ratings for men at the finish line (164 km) were as follows: thermal, 2.5-7.5; thirst, 1-8; perceived exertion, 12-19; pain, 0-7; and muscle cramp, 0-4.

Table 5 presents energy expenditure values for the Event Day. These calculations involve the 9-hour cycling event, energy expenditure during the remainder of the day (15 h of sleep + activities), and the sum of these two quantities. Based on this total daily energy expenditure (column 4), metabolic water production was calculated as an adjunct to the body mass change data (Table 2) and the fluid consumption data (Table 6).

Tables 6 and 7 present dietary intakes of male and female cyclists, on the day before the event (Day -1) and on Event Day. These values include the total dietary fluid volume, energy (kcal), and selected macronutrients. Figure 4 depicts the types of fluids that were consumed by men on Day -1 and on Event Day.

## **DISCUSSION**

This 164-km cycling event, conducted in Texas during the month of August, proved to be challenging for cyclists because the ambient temperature, with minimal cloud cover, exceeded 39.0°C during the final 2 h of exercise. This event also proved to be logistically challenging for investigators at distant data measurements sites because (a) it was difficult to identify test participants among thousands of riders, and (b) the temporal difference between the first and last test participant was 4.2 h, a road distance of approximately 61 km. Interestingly, the mean physiological and perceptual

responses of women and men were similar, except the variables that were related to body size and energy-metabolic requirements. Although we did not test for significant differences between men and women, due to the small number of women who enrolled (n=6), we include women's data because few previous studies have examined competitive female cyclists (9).

### **Physiological and Perceptual Responses**

We initially hypothesized that cyclists would not maintain thermal, fluid and energy balance, and that physiological measures would indicate great stress. However, the average completion time was 9.0 – 9.1 h, and cyclists maintained an overall ground speed of 17.8–17.9 km·h<sup>-1</sup> (Table 1), the mean values indicated mild-to-moderate thermal and cardiovascular strain. Also interesting are the mean values for body mass change, urine specific gravity, urine color, serum osmolality, and serum electrolytes (Tables 2 and 3), which show that cyclists maintained fluid-electrolyte balance during this event. But there were exceptions. Those who responded at the extremes of the frequency distributions (Figures 1-3) were the cyclists whose homeostasis was perturbed most and who had the greatest risk of exertional heat illness. For example, the two greatest T<sub>GI</sub> values at the finish line were 39.4 and 40.2°C (Fig. 1), meaning that 13% (2/15) of the cyclists in this subsample were hyperthermic, at the threshold of exertional heatstroke (2), but had no signs or symptoms of heat illness. Eleven participants exhibited urine specific gravity values ≥1.030, indicating frank dehydration (28). In fact, the six greatest values for urine specific gravity (range, 1.034 - 1.038; Fig. 2) were extraordinarily high, well outside the 95% confidence limits published previously for young men during daily activities (4). The frequency distribution in Figure 3 also is

interesting because it demonstrates that four cyclists lost more than 4% body mass during this event. This level of dehydration is widely recognized as one that degrades endurance exercise performance (28).

Numerous studies have been published regarding renal function after a marathon and cross-country skiing but, because little is known about the physiological effects or fluid-electrolyte balance during ultra-endurance cycling (23), we present the following findings regarding renal function and fluid-electrolyte balance. First, sweat rate on Event Day was  $1.13 \pm 0.54 \text{ L} \cdot \text{h}^{-1}$ , in a subset of 20 men during the first 52 km. Extrapolating this sweat loss to 164 km, we calculated that the total mean sweat loss was 3.56 L. Second, the male cyclists consumed 5.91 L of fluid during this event (Table 6). Third, because we were unable to measure the mass of consumed solid food, urine volume, and fecal mass, we assumed that the solid food mass gain and the fecal mass loss were approximately equal (0.5 kg) across 9 h (7), and assumed that respiratory water loss and metabolic water production (0.35 kg; Table 5) were approximately equal (1). Considering a net body mass change of -0.76 kg (Table 2), we then calculated that mean urine excretion during exercise was 3.1 L ( $-3.56 + 5.91 - \text{urine} = -0.76$ ). This rate of urine production ( $\sim 300 \text{ ml} \cdot \text{h}^{-1}$ ) suggests that 9 h of exercise in the heat of Texas did not pose a threat to renal health, in agreement with the findings of Neumayr et al. (23), who concluded that ultramarathon cycling was “harmless” to the renal function of 16 men who raced across the Alps in 26.4 h, in ambient temperatures ranging from -1 to 20°C. Yet, although an average of 0.8% of body mass was lost during the HHH (Table 2), Figure 2 suggests that considerable renal concentrating stress existed in 26% of

male cyclists, in that they experienced a urine specific gravity  $\geq 1.030$ . This phenomenon, resulting from exercise-heat stress, deserves further study.

**Perceptual Ratings.** Although little is known about changes that occur in athlete perceptions during ultra-endurance cycling in a hot environment, the ratings in Table 4 provide several insights about non-elite cyclists. As hypothesized, perceptual ratings increased significantly ( $P < .025$  to  $.001$ ) from the start (0 km) to the finish (164 km). However, although the mean values indicated only mild-to-moderate strain, the range of values was large for each rating. For example, thermal stress at the finish line ranged from “cool” to “very hot”; thirst, from “not thirsty at all” to “very thirsty”; perceived exertion, from “somewhat hard” to “very, very hard”; and pain intensity, from “no pain at all” to “very strong pain”. In combination with cumulative ground speed (Table 1), the perceptual ratings in Table 4 provide a consistent picture of the HHH experience of non-elite cyclists. That is, as their ground speed declined between aid stations (0-52 km, 52-97 km, 97-164 km), they encountered greater pain and muscle cramps; they also felt hotter, thirstier, and more fatigued. Every rider covered the final stage of the course (97-164 km) slower than the previous two stages (0-52 km, 52-97 km) due, in part, to an air temperature exceeding  $39^{\circ}\text{C}$  during the final 2 h of the event.

### **Nutritional Considerations**

**Energy Balance.** Observations of professional cyclists during the Tour de France (26) indicated that their average daily energy expenditure exceeded  $5,736 \text{ Kcal}\cdot 24\text{h}^{-1}$  ( $24 \text{ MJ}\cdot 24\text{h}^{-1}$ ). Utilizing methods that were developed by other research teams (17,26,29), Table 5 describes the following components of energy expenditure on Event Day: cycling (9 h), sleep + non-exercise daily activities (15 h), and 24 h total

energy expenditure. The data for 33 men show that 64.1% ( $2950/4602 = 0.641$ ) of energy expenditure on Event Day was expended during 9 h of cycling exercise, on the 164-km course. The average energy expenditure for men on Event Day was 328 Kcal·h<sup>-1</sup> during 9 h of cycling, and 110 Kcal·h<sup>-1</sup> during 15 h of non-exercising daily activities; the former was almost three times greater than the latter. During the event, the energy expended (at a mean ground speed of 17.9 km·h<sup>-1</sup>) was equivalent to 18.3 Kcal·km<sup>-1</sup> (29.5 Kcal·mi<sup>-1</sup>). Assuming that these cyclists (Table 5) awoke at 0530 h, and began exercise at 0700 h (race start time), we calculated that their energy expenditure for all activities was 3,115 Kcal (13.0 MJ) up to the race finish at 1600 h. In comparison, these men consumed only 521 Kcal (2.2 MJ) up to 1600 h (Table 6), in fluids and solid foods. This means that they consumed only 16.7% of the energy they expended (i.e., an energy deficit of 2,594 Kcal, 10.9 MJ), supporting the previously-described phenomenon of voluntary energy depletion during endurance cycling (7). Brouns et al. (8) provided two theoretical explanations for such voluntary energy depletion: (a) the amount of feces produced from a normal diet would require competitors to defecate several times during the day, consequently interrupting exercise; and (b) a large volume of solid food may interfere with fluid intake, which is important on hot days to replace large sweat losses. In a related study (7), they concluded that cyclists may not be able to eat enough food to compensate for the extremely high energy expenditure of ultra-endurance competition, even though they are well aware of the importance of adequate energy intake.

Although we have no data to indicate which foods or fluids were consumed after the HHH event (Tables 6 and 7), Brouns et al. (7,8) provide a plausible scenario. They

observed the dietary intakes of 13 highly trained males (age,  $20 \pm 2$  yr; body mass,  $73.3 \pm 1.7$  kg; height,  $180 \pm 2$  cm) who consumed a diet high in carbohydrates on two baseline days. On two subsequent days of simulated Tour de France competition ( $50 - 90\% \text{VO}_{2\text{max}}$ , intermittent,  $6.5 \text{ h}\cdot\text{d}^{-1}$ ), these cyclists experienced energy deficits of 10 and  $8 \text{ MJ}\cdot 24\text{h}^{-1}$ . On the single recovery day, which followed these prolonged exercise simulations, investigators measured a mean excess energy intake of  $\sim 2 \text{ MJ}\cdot 24 \text{ h}^{-1}$  and concluded that energy intake increased in proportion to the energy deficit.

**Macronutrients.** Carbohydrate intake represents the most striking feature of the macronutrient analysis presented in Table 7. Eighty-four percent of dietary energy on Event Day (versus 55.2% on Day -1) was consumed in the form of carbohydrates. However, all other macronutrients were considerably under-consumed by men on Event Day. For example, the large percentage of dietary carbohydrate on Event Day was consumed at the expense of protein (Table 7). Mean protein consumption between 0530 – 1600 h was only  $0.1 \text{ g}\cdot\text{kg body mass}^{-1}$  (8 g), which was well below the  $1.2 - 1.4 \text{ g}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$  recommended by Lemon (20) and Tarnopolsky et al. (30), but similar to that of male professional cyclists during the Tour of Spain ( $167 \text{ km}\cdot\text{d}^{-1}$ , two mountain stages; (15). Considering that post-exercise protein intake has been associated with reduced post-exercise muscle soreness (14) and decreased post-exercise skeletal muscle damage (27), we recommend that dietitians and sport nutritionists counsel recreational cyclists to consume protein preferentially, after ultra-endurance events such as the HHH. Our recommendation is supported by the suggestion that elite competitors have better nutrition knowledge, and superior on-course nutritional provisions, than slower recreational cyclists (25).

Two other observations regarding carbohydrate are noteworthy. Firstly, Coyle (12) and the American College of Sports Medicine (28) recommend that cyclists consume 30-60 g carbohydrate during competition, to maintain blood glucose and sustain exercise performance. If cyclists had complied with this recommendation, their carbohydrate intake would have been 270 – 540 g on Event Day. The fact that no cyclist consumed this amount of carbohydrate (range of intakes, 17 – 207 g·10.5 h<sup>-1</sup>; Table 7) may explain, in part, why the ground speed decreased during each successive segment (0 to 52 km, 29.7 ± 4.3 km·h<sup>-1</sup>; 52 to 97 km, 24.1 ± 4.1 km·h<sup>-1</sup>; 97 to 164 km, 12.7 ± 1.8 km·h<sup>-1</sup>; mean ± SD). Secondly, only 43% of fluids consumed on Event Day (Fig. 4) were carbohydrate-electrolyte replacement beverages (C-E), and 49% were water. This low C-E intake by cyclists does not follow the widely-accepted sport nutrition principles that (a) hypoglycemia influences central fatigue (6), (b) endurance is enhanced by pre-exercise glycogen levels (19), and (c) exhaustion from hypoglycemia (6) and glycogen depletion (19) can be postponed by carbohydrate feeding during exercise. Also contrary to these principles, the cyclist who consumed the smallest amount of carbohydrate on Event Day (17 g in 10.5 h) crossed the finish line 1 h ahead of the closest male study participant (range of 41 other men, 17 – 207 g protein; Table 7). He also consumed little food before and during the event (82 Kcal total, 81% carbohydrate), as opposed to Day -1 (2074 Kcal, 69% carbohydrate). We interpret this case to mean that carbohydrate intake is only one factor that influences exercise performance. This is acknowledged as a limitation of the present study, and may explain why the statistical correlation between carbohydrate intake (Event Day, n = 27)

and finish time was weak and non-significant, for the 164-km total time ( $r^2 = .02$ ;  $P = .44$ ) and for the time to complete the final 97 km ( $r^2 = .02$ ;  $P = .49$ ).

**Fluids and Sodium.** The mean volume of fluid which men consumed on Day -1 was  $2.17 \pm 1.64 \text{ L}\cdot 24 \text{ h}^{-1}$  (Table 6)--considerably different from 10 male professional cyclists (15). The latter consumed only an average of 1.26 L of fluid during 3 d of competition; water comprised 82% of their total fluid volume while covering 167 km·d<sup>-1</sup>. However, the mean volume of fluid consumed on Event Day (Table 6;  $5.91 \pm 2.38 \text{ L}\cdot 10.5 \text{ h}^{-1}$ ) was 2.7 times greater than Day -1. This is reasonable, considering the 3.56 L sweat loss (see above) which men experienced during the 164-km ride, and explains maintenance of body mass throughout the HHH (Table 2). The fluid intake of women on Event Day, expressed per kilogram of body mass, was nearly identical to men ( $71 \text{ ml}\cdot\text{kg}^{-1}$ , Table 6). This supports observations of marathon runners (16) in which 54 women drank virtually the same amount of water as 63 men (22.9 and 22.3 cups, respectively) during a 42.1 km footrace.

It also is important to note that the fluids which men consumed on Day -1 (Fig. 4) consisted of eight varieties, whereas C-E and water comprised 92% of the fluids consumed on Event Day. Figure 4, which shows the ratio of water (49%) to C-E beverages (43%) consumed during the HHH event, stands in contrast to the findings of García-Rovés et al. (15), who reported that 10 elite competitors drank considerably more water (82%) than C-E (18%) during three consecutive days of the Tour of Spain. This may be due to the greater consumption of solid foods by elite cyclists. These findings (i.e., regarding relatively large percentages of water) deserve further study,

since numerous experiments have reported that blood glucose concentration is lower during water (control) experiments than during C-E trials (6).

In recent years, it has become customary for race directors to provide sodium supplements (i.e., salt capsules) at the aid stations of ultra-endurance events (24). This practice is based, in part, on the importance of sodium in the maintenance of extracellular volume, as well as the relationship between whole-body sodium imbalance, heat exhaustion, and heat cramps (2). Some sport nutrition authorities recommend that athletes who compete in events longer than 6-8 h should consume sodium at a rate of  $\sim 1000 \text{ mg}\cdot\text{h}^{-1}$ , in either C-E or foods (10). The data in Table 7 demonstrate that the mean dietary sodium intake of men on Event Day was low ( $852 \pm 531 \text{ mg}$  total), only 19% of Day -1. However, despite a total sweat loss of 3.56 L (see above), we have no evidence to indicate that this deficit was detrimental to health or performance. For example, the mean serum sodium level at the finish line (Table 3,  $141 \pm 3 \text{ mEq}\cdot\text{L}^{-1}$ ) was well within the normal range. Secondly, the rating of muscle cramp severity (Table 4) at the finish line was weakly and non-significantly correlated with the amount of sodium consumed (Table 7;  $r^2 = .03$ ), the total volume of fluid consumed (all varieties; Table 6;  $r^2 = .02$ ), and the total volume of water consumed ( $r^2 = .01$ ). We interpret these observations to mean that the whole-body sodium imbalance was not great enough to result in clinical signs or symptoms of heat cramps or heat exhaustion (2).

## **PRACTICAL APPLICATIONS**

The participants in this study were considered to be non-elite cyclists, on the basis of their average performance time (9.0 and 9.1 h) and ground speed (17.9 and

17.8 km·h<sup>-1</sup>). We recommend that dietitians, sport nutritionists and clinicians counsel recreational ultra-endurance cyclists to emphasize the following:

1. Unrecognized under-consumption of energy occurs. In the present 100 km event, an energy deficit of 2,594 Kcal, (10.9 MJ) was measured. Cyclists should consume considerably more energy during the event, to maintain exercise performance (12).
2. Protein is under-consumed on Event Day. During 2-4 post-event meals, protein should be emphasized by consuming at least 1.2 – 1.4 g per kg body mass.

Further, we remind clinicians, coaches and athletes that a minority of cyclists in ultra-endurance events experience extreme final values (i.e., urine specific gravity of 1.035 to 1.038, body mass loss >4 kg, and T<sub>GI</sub> of 39.4 and 40.2). These levels of renal and thermal stress re-emphasize the widely-recognized principles of (a) maintaining adequate fluid intake to avoid severe dehydration, as guided by multiple observations of urine color or urine specific gravity (4,28); (b) learning and watching for the signs and symptoms of the exertional heat exhaustion and exertional heatstroke (2); and (c) participating with a companion who will share the responsibility of monitoring health and nutrition, during and after the event (2,10).

## REFERENCES

1. Armstrong, LE, and Casa, DJ. Methods to evaluate electrolyte and water turnover of athletes. *Athl Train Sport Hlth Care* 1: 169-179, 2009.

2. Armstrong, LE, Casa, DJ, Millard-Stafford, M, Moran, DS, Pyne, SW, and Roberts, WO. Exertional Heat Illness during Training and Competition. American College of Sports Medicine Position Stand. *Med Sci Sports Exerc* 39(3): 556-572, 2007.
3. Armstrong, LE, Maresh, CM, Castellani, J, Bergeron, M, Kenefick, RW, LaGasse, KE, and Riebe D. Urinary indices of hydration status. *Intl J Sport Nutr* 4: 265-279, 1994.
4. Armstrong, LE, Pumerantz, AC, Fiala, KA, Roti, MW, Kavouras, SA, Casa, DJ, and Maresh, CM. Human Hydration Indices: Acute and Longitudinal Reference Values. *Intl J Sport Nutr Exerc Metab* 20: 145-153, 2010.
5. Borg, G. Perceived exertion as an indicator of somatic stress. *Scand J Rehabil Med* 2: 92-98, 1970.
6. Brouns, F, Rehrer, NJ, Saris, WHM, Beckers, E, Menheere, P, and ten Hoor, F. Effect of carbohydrate intake during warming-up on the regulation of blood glucose during exercise. *Int J Sports Med* 10(suppl 1): S68-S75, 1989a.
7. Brouns, F, Saris, WHM, Stroecken, J, Beckers, E, Thijssen, R, Rehrer, NJ, and ten Hoor F. Eating, drinking, and cycling. A controlled Tour de France simulation study, Part I. *Int J Sports Med* 10: Suppl 1, S32-S40, 1989b.
8. Brouns, F, Saris, WHM, Stroecken, J, Beckers, E, Thijssen, R, Rehrer, NJ, and ten Hoor F. Eating, drinking, and cycling. A controlled Tour de France simulation study, Part II. Effect of diet manipulation. *Int J Sports Medicine* 10: Suppl 1, S41-S48, 1989c.
9. Burke, LM. Nutritional Practices of Male and Female Endurance Cyclists. *Sports Med* 31(7): 521-532, 2001.

10. Clark, N, Tobin, J, and Ellis, C. Feeding the ultraendurance athlete: Practical tips and a case study. *J Am Diet Assoc* 92: 1258-1262, 1992
11. Cook, DB, O'Connor, PJ, Eubanks, SA, Smith, JC, and Lee, M. Naturally occurring muscle pain during exercise: assessment and experimental evidence. *Med Sci Sport Exerc* 29: 999-1012, 1997.
12. Coyle, E. Fluid and fuel intake during exercise. *J Sport Sci* 22: 39-55, 2004.
13. Engell, DB, Maller, O, Sawka, MN, Francesconi, RN, Drolet, L, and Young, AJ. Thirst and fluid intake following graded hypohydration levels in humans. *Physiol Behav* 40: 229-236, 1987.
14. Flakoll, PJ, Judy, T, Flinn, K, Carr, C, and Flinn, S. Postexercise protein supplementation improves health and muscle soreness during basic military training in marine recruits. *J Appl Physiol* 96: 951-956, 2004.
15. García-Rovés, PM, Terrados, N, Fernández, SF, and Patterson, AM. Macronutrients intake of top level cyclists during continuous competition—Change in the feeding pattern. *Int J Sports Med* 19: 61-67, 1998.
16. Hew, TD. Women hydrate more than men during a marathon race. *Clin J Sports Med* 15: 148-153, 2005.
17. Hoyt, RW, and Honig, A. Environmental influences on body fluid balance during exercise: Altitude. In: *Body Fluid Balance: Exercise and Sport*. Buskirk, ER, and Puhl, SM, eds. Boca Raton, FL: CRC Press, 1996. pp. 183–196.
18. Kratz, A, Ferraro, M, Sluss, PM, and Lewandrowski, KM. Case records of the Massachusetts General Hospital. Laboratory reference values. *N Engl J Med* 351: 1548-1563, 2004.

19. Kuipers, H, Saris, WH, Brouns, F, Keizer, HA, and ten Bosch C. Glycogen synthesis during exercise and rest with carbohydrate feeding in males and females. *Int J Sports Med* 10(suppl 1): S63-S67, 1989.
20. Lemon, PWR. Dietary protein requirements in athletes. *J Nutr Biochem* 8: 52-60, 1997.
21. Maughan, RJ. Thermoregulation in marathon competition at low ambient temperature. *Int J Sports Med* 6(1): 15-19, 1985.
22. Maughan, RJ, Leiper, JB, and Thompson J. Rectal temperature after marathon running. *Brit J Sports Med* 19(4): 192-196, 1985.
23. Neumayr, G, Pfister, R, Hoertnagl, H, Mitterbauer, G, Prokop, W, and Joannidis, M. Renal function and plasma volume following ultramarathon cycling. *Int J Sports Med* 26(1): 2-8, 2005.
24. Noakes, T. Hyponatremia in Distance Runners: Fluid and Sodium Balance During Exercise. *Curr Sports Med Rep* 4: 197–207, 2002.
25. Robins, A, and Hetherington, MM. A comparison of pre-competition eating patterns in a group of non-elite triathletes. *Intl J Sport Nutr Exerc Metab* 15: 442-457, 2005.
26. Saris, WHM, van Erp-Baart, MA, Brouns, F, Westerterp, KR, and ten Hoor, F. Study on food intake and energy expenditure during extreme sustained exercise: The Tour de France. *Int J Sports Med* 10(suppl 1): S26-S31, 1989.
27. Saunders, M, Kane, M, and Todd, K. Effects of a carbohydrate–protein beverage on cycling endurance and muscle damage. *Med Sci Sports Exerc* 36: 1233–1238, 2004.

28. Sawka, MN, Burke, LM, Eichner, ER, Maughan, RJ, Montain, SJ, and Stachenfeld, NS. Exercise and fluid replacement. American College of Sports Medicine Position Stand. *Med Sci Sports Exerc* 39(2): 377-90, 2007.
29. Swain, DP, Coast, JR, Clifford, PS, Milliken, MC, and Stray-Gundersen, J. Influence of body size on oxygen consumption during bicycling. *J Appl Physiol* 62(2): 668-72, 1987.
30. Tarnopolsky, MA, MacDougall, JD, and Atkinson, SA. Influence of protein intake and training status on nitrogen balance and lean body mass. *J Appl Physiol* 64: 187-193, 1988.
31. USA Cycling. USA Cycling fact sheet. Colorado Springs, CO. 2010.
32. Young, AJ, Sawka, MN, Epstein, Y, Decristofano, B, and Pandolf, KB. Cooling different body surfaces during upper and lower body exercise. *J Appl Physiol* 63: 1218-1223, 1987.

## **ACKNOWLEDGEMENTS**

Keith H. Williamson, M.D. is the medical director of the HHH. The authors acknowledge the technical skills of Lynn Mardon and Michael Judd. Present affiliations of the authors include: M.S. Ganio, University of Arkansas, Fayetteville, Arkansas; E.C. Lee, Mt Desert Island Biological Laboratory, Salisbury Cove, ME; B.P. McDermott, University of Tennessee, Chattanooga, TN ; J.L. Vingren, University of North Texas, Denton, TX; and J.E. Wingo, University of Alabama, Tuscaloosa, AL.

## Figure Legends

Figure 1. Gastrointestinal temperature ( $T_{GI}$ ) at the end of the 164-km cycling event, measured in a subset of participants.

Figure 2. Distribution of urine specific gravity values for men and women, measured at the end of 9.0 – 9.1 h of exercise.

Figure 3. Frequency distribution of percent change of body mass, from pre-exercise (0 km) to post-exercise (164 km).

Figure 4. Types of fluids consumed by men ( $n = 31$ ). On Day -1, the other category consisted of 5% fruit juice, 3% pickle juice, and 2% milk. On Event Day, the other category consisted of 6% pickle juice, and 1% milk. Abbreviations: C-E, carbohydrate-electrolyte replacement fluid; CSD, carbonated soft drink.

**Table 1. Mean ( $\pm$  SD) elapsed time, gastrointestinal temperature and heart rate measured before the start, at two aid stations, and at the finish line.**

<b>Variable</b>	<b>Group (n)</b>	<b>Pre-Race (0 km)</b>	<b>Aid Station (52 km)</b>	<b>Aid Station (97 km)</b>	<b>Finish (164 km)</b>
<b>Elapsed Time (h) <sup>a</sup></b>	Men (n = 33)	0.0	1.8 $\pm$ 0.3	3.7 $\pm$ 0.5	9.1 $\pm$ 1.2
	Women (n = 5) <sup>b</sup>	0.0	1.8 $\pm$ 0.2	3.8 $\pm$ 0.2	9.0 $\pm$ 0.2
<b>Cumulative Ground Speed (km·h<sup>-1</sup>)</b>	Men (n = 33)	0.0	29.5 $\pm$ 4.7	26.6 $\pm$ 3.8 <sup>c</sup>	17.9 $\pm$ 2.4 <sup>d</sup>
	Women (n = 5) <sup>b</sup>	0.0	29.3 $\pm$ 3.7	25.8 $\pm$ 1.2	17.8 $\pm$ 0.4
<b>Gastrointestinal Temperature (°C)</b>	Men (n = 12)	36.86 $\pm$ 0.38	38.14 $\pm$ 0.46 <sup>e</sup>	38.16 $\pm$ 0.21 <sup>e</sup>	38.50 $\pm$ 0.39 <sup>f</sup>
	Women (n = 3) <sup>b</sup>	37.02 $\pm$ 0.09	38.26 $\pm$ 0.32	38.24 $\pm$ 0.52	38.10 $\pm$ 0.28
<b>Heart Rate (beats·min<sup>-1</sup>)</b>	Men (n = 16)	<sup>g</sup>	149 $\pm$ 9	155 $\pm$ 7 <sup>c</sup>	161 $\pm$ 12 <sup>d</sup>
	Women (n = 3) <sup>b</sup>	<sup>g</sup>	140 $\pm$ 24	151 $\pm$ 12	149 $\pm$ 14

<sup>a</sup>, subjects stopped for approximately 5-10 min at two aid stations for research measurements, elimination, drinking, and eating

<sup>b</sup>, no statistical comparison was performed across time for women, due to small sample size and incomplete data sets; this applies to all tables and figures below

<sup>c</sup>, significantly different from 52 km ( $P < .001$ ) and 164 km ( $P < .025$ ) values for men

<sup>d</sup>, significantly different from 52 km ( $P < .001$ ) values for men

<sup>e</sup>, significantly different from 0 km ( $P < .001$ ) values for men

<sup>f</sup>, significantly different ( $P < .05$  to  $.001$ ) from all other means for men

<sup>g</sup>, not measured

**Table 2. Mean ( $\pm$  SD) changes of body mass and two urinary indices of hydration for men (n = 42) and women (n = 6), measured at four locations.**

<b>Variable</b>	<b>Group</b>	<b>Pre-Race (0 km)</b>	<b>Aid Station (52 km)</b>	<b>Aid Station (97 km)</b>	<b>Finish (164 km)</b>
<b>Body Mass (kg)</b>	Men	85.85 $\pm$ 14.79	84.82 $\pm$ 14.55	85.77 $\pm$ 14.78	85.09 $\pm$ 14.47
	Women	67.32 $\pm$ 7.21	66.31 $\pm$ 7.06	67.48 $\pm$ 7.49	67.25 $\pm$ 7.48
<b>Body Mass Change (kg)</b>	Men	0.0	- 1.03 $\pm$ 0.98	- 0.08 $\pm$ 1.27	- 0.76 $\pm$ 1.82
	Women	0.0	- 1.00 $\pm$ 0.54	+ 0.17 $\pm$ 0.79	- 0.07 $\pm$ 0.94
<b>Urine Specific Gravity</b>	Men	1.019 $\pm$ 0.008	1.016 $\pm$ 0.010	1.018 $\pm$ 0.010	1.023 $\pm$ 0.008 <sup>a</sup>
	Women	1.018 $\pm$ 0.001	1.017 $\pm$ 0.006	1.022 $\pm$ 0.005	1.023 $\pm$ 0.004
<b>Urine Color <sup>b</sup></b>	Men	4 $\pm$ 2	4 $\pm$ 2	5 $\pm$ 2 <sup>b</sup>	6 $\pm$ 1 <sup>c</sup>
	Women	4 $\pm$ 1	5 $\pm$ 1	5 $\pm$ 2	6 $\pm$ 1

<sup>a</sup>, significantly greater (P<.001) than all other time points for men, as determined by ANOVA

<sup>b</sup>, significantly greater (P<.05) than 0 km for men

<sup>c</sup>, significantly greater (P<.001) than all other time points for men

**Table 3. Serum osmolality and electrolytes (mean  $\pm$  SD) measured before and after exercise.**

<b>Constituent</b>	<b>Group <sup>a</sup></b>	<b>Pre-Exercise (0 km)</b>	<b>Finish (164 km)</b>
<b>Osmolality (mOsm·kg<sup>-1</sup>)</b>	Men	291 $\pm$ 4	292 $\pm$ 6
	Women	291 $\pm$ 3	289 $\pm$ 4
<b>Sodium (mEq·L<sup>-1</sup>)</b>	Men	141 $\pm$ 1	141 $\pm$ 3
	Women	141 $\pm$ 1	140 $\pm$ 3
<b>Chloride (mEq·L<sup>-1</sup>)</b>	Men	102 $\pm$ 2	102 $\pm$ 4
	Women	104 $\pm$ 1	104 $\pm$ 4
<b>Potassium (mEq·L<sup>-1</sup>)</b>	Men	4.1 $\pm$ 0.3	4.5 $\pm$ 0.5 <sup>b</sup>
	Women	4.4 $\pm$ 0.5	5.0 $\pm$ 0.5

<sup>a</sup>, blood was sampled from 33 men and 5 women

<sup>b</sup>, significantly greater ( $P < .001$ ) than Pre-Exercise for men, as determined by Student's t-test

**Table 4. Perceptual ratings <sup>a</sup> (mean ± SD) for male (n = 40) and female (n = 6) cyclists, during the 164 km mass participation event.**

<b>Sensation</b>	<b>Group</b>	<b>Pre-Exercise (0 km)</b>	<b>Aid Station (52 km)</b>	<b>Aid Station (97 km)</b>	<b>Finish (164 km)</b>
<b>Thermal</b>	Men	4.0 ± 0.6	4.7 ± 0.6 <sup>b</sup>	5.2 ± 0.7 <sup>b,c</sup>	5.8 ± 1.0 <sup>d</sup>
	Women	4.1 ± 0.2	4.1 ± 0.2	5.2 ± 0.3	5.5 ± 1.1
<b>Thirst</b>	Men	2.2 ± 1.1	2.9 ± 1.1 <sup>b</sup>	3.6 ± 1.4 <sup>b,c</sup>	5.0 ± 1.7 <sup>d</sup>
	Women	3.8 ± 1.3	3.8 ± 1.7	4.7 ± 1.6	5.8 ± 1.8
<b>Perceived Exertion</b>	Men	e	13 ± 2	14 ± 1 <sup>c</sup>	15 ± 2 <sup>d</sup>
	Women	e	12 ± 1	13 ± 2	14 ± 4
<b>Pain</b>	Men	0.1 ± 0.2	0.5 ± 0.8 <sup>b</sup>	1.6 ± 1.1 <sup>b,c</sup>	2.6 ± 1.5 <sup>d</sup>
	Women	0.0 ± 0.0	0.1 ± 0.2	1.5 ± 1.4	1.8 ± 1.9
<b>Muscle cramp</b>	Men	0.0 ± 0.2	0.2 ± 0.4	0.4 ± 0.6 <sup>b,c</sup>	1.2 ± 1.3 <sup>d</sup>
	Women	0.0 ± 0.0	0.0 ± 0.0	0.3 ± 0.5	0.7 ± 0.8

<sup>a</sup>, perceptual rating scales are described in the Methods section above; the numerical ranges for each measurement appear in the Results section above

<sup>b</sup>, significantly greater (P<.001) than 0 km for men, as determined by ANOVA

<sup>c</sup>, significantly greater (P<.025 to .001) than 52 km for men

<sup>d</sup>, significantly greater (P<.001) than all other time points for men

<sup>e</sup>, rated only at aid stations and the finish line

**Table 5. Calculated energy expenditure and metabolic water production (mean  $\pm$  SD) of men (n = 33) and women (n = 5) on Event Day.**

Cyclists	9h Energy Expenditure (Cycling) <sup>a</sup>		15h Energy Expenditure (Sleep + Activities) <sup>b</sup>		24h Total Energy Expenditure <sup>c</sup>		9h (Cycling) Metabolic Water Production (ml) <sup>d</sup>	24h Total Metabolic Water Production (ml) <sup>d</sup>
	MJ	Kcal	MJ	Kcal	MJ	Kcal		
Men	12.3 $\pm$ 1.8	2950 $\pm$ 436	6.9 $\pm$ 1.0	1652 $\pm$ 248	19.2 $\pm$ 2.3	4602 $\pm$ 538	349 $\pm$ 52	545 $\pm$ 64
Women	11.6 $\pm$ 0.5	2764 $\pm$ 123	5.6 $\pm$ 0.6	1328 $\pm$ 149	17.1 $\pm$ 1.1	4092 $\pm$ 266	327 $\pm$ 15	485 $\pm$ 32

<sup>a</sup>, column 2 calculations utilized the over-ground cycling method of Swain et al.(29). Oxygen consumption was calculated from the ground speed and body mass of each rider, then was converted to energy expenditure (Kcal = L·min<sup>-1</sup> x 5) by considering total exercise time of each subject.

<sup>b</sup>, column 3 calculations used the method of Saris et al. (26). 15 h daily resting energy expenditure = 7h sleep (estimated as 0.074 KJ/min/kg) + 8h of other non-exercise activities (estimated as 0.111 KJ/min/kg).

<sup>c</sup>, column 4 = column 2 + column 3

<sup>d</sup>, column 5 and 6 calculations involved the method of Hoyt & Honig (17):metabolic water (ml) = (0.119 x EE) – 2.25, where EE is energy expenditure

**Table 6. Total energy and fluid intake (mean  $\pm$  SD) of cyclists on Day -1 and Event Day. <sup>a</sup>**

Variable	Group <sup>b</sup>	Units	Day Before Event (Day -1) <sup>c</sup>		Event Day <sup>c</sup>	
			Mean $\pm$ SD	Range	Mean $\pm$ SD	Range
		MJ	11.1 $\pm$ 3.5	7.2 – 21.4	2.2 $\pm$ 0.9	0.4 – 4.1
		Kcal	2648 $\pm$ 837	581 – 5049	521 $\pm$ 224	84 – 967
		MJ	7.8 $\pm$ 2.7	5.6 – 11.4	2.4 $\pm$ 1.5	1.0 – 4.7
		Kcal	1865 $\pm$ 639	1352 - 2725	570 $\pm$ 347	247 – 1123
		KJ·kg <sup>-1</sup>	132.7 $\pm$ 46.0	34.6 – 249.6	26.6 $\pm$ 12.0	4.1 – 52.4
		kcal·kg <sup>-1</sup>	31.7 $\pm$ 11.0	8.3 – 59.6	6.4 $\pm$ 2.9	1.0 – 12.0
		KJ·kg <sup>-1</sup>	115.0 $\pm$ 34.0	84.2 – 168.1	35.9 $\pm$ 22.3	17.0 – 69.3
		kcal·kg <sup>-1</sup>	27.5 $\pm$ 8.1	20.1 – 40.1	8.6 $\pm$ 5.3	4.1 – 16.5
Fluid Consumed in all Beverages <sup>d</sup>	Men	L	2.17 $\pm$ 1.64	0.33 – 6.03	5.91 $\pm$ 2.38	1.89 – 10.88
	Women	L	1.28 $\pm$ 1.53	0.12 – 4.02	4.68 $\pm$ 1.66	2.78 - 6.83
Fluid Consumed in all Beverages <sup>d</sup>	Men	ml·kg <sup>-1</sup>	25.6 $\pm$ 19.2	5.0 – 78.2	70.6 $\pm$ 28.1	22.1 – 138.0
	Women	ml·kg <sup>-1</sup>	19.1 $\pm$ 22.7	1.9 – 60.5	71.1 $\pm$ 29.6	38.2 – 117.8

<sup>a</sup>, consumed in fluids and solid foods

<sup>b</sup>, men, n = 33; women, n = 5

<sup>c</sup>, Day -1 occurred on Friday and the event occurred on Saturday; values for Day -1 represent 24h; Event Day values represent the morning meal plus food intake during the event (~10.5h), but include no food after the event; thus, no between-day statistical comparisons were made

<sup>d</sup>, fluid varieties appear in Figure 4

**Table 7. Consumption of selected macronutrients by men (n = 33) and women (n = 5) on Day -1 and Event Day.**

Macronutrient <sup>a</sup>	Group	Day Before Event (Day -1) <sup>b</sup>		Event Day <sup>b</sup>	
		Mean ± SD	Range	Mean ± SD	Range
<b>Carbohydrate (g)</b>	Men	368 ± 130	116 – 722	106 ± 48	17 – 207
	Women	236 ± 112	135 – 304	116 ± 80	52 - 243
<b>Carbohydrate (% of total MJ)</b>	Men	56.7 ± 11.4	33.4 – 79.6	81.6 ± 12.3	47.8 – 100.0
	Women	49.3 ± 9.5	40.1 – 64.6	79.1 ± 10.4	61.2 – 86.5
<b>Protein (g)</b>	Men	93 ± 40	46 – 167	8 ± 7	0 – 29
	Women	92 ± 34	64 – 139	7 ± 3	3 – 12
<b>Protein (g·kg body mass<sup>-1</sup>)</b>	Men	1.1 ± 0.5	0.4 – 2.1	0.1 ± 0.1	0.0 – 0.3
	Women	1.4 ± 0.4	0.9 – 2.1	0.1 ± 0.1	0.1 – 0.2
<b>Protein (% of total MJ)</b>	Men	13.8 ± 3.7	8.2 – 22.7	6.6 ± 4.3	0.4 – 16.9
	Women	19.6 ± 1.3	17.5 – 20.6	5.9 ± 3.2	3.3 – 11.3
<b>Sodium (mg)</b>	Men	4475 ± 2072	1156 - 11285	852 ± 531	124 - 2352
	Women	3453 ± 2134	1809 - 6351	911 ± 562	413 -1940
<b>Potassium (mg)</b>	Men	2255 ± 1256	535 - 5668	434 ± 221	59 - 935
	Women	2292 ± 1255	1297 - 4774	442 ± 240	149 - 773

<sup>a</sup>, consumed in fluids and sold food

<sup>b</sup>, see footnote c from Table 6